

FIXED POINT STRUCTURE OF PADÉ-SUMMATION APPROXIMATIONS TO THE QCD β -FUNCTION

V. Elias*, F. Chishtie

Department of Applied Mathematics, The University of Western Ontario, London, Ontario N6A 5B7, CANADA

**E-mail: zohar@apmaths.uwo.ca*

T.G. Steele

Department of Physics and Engineering Physics, University of Saskatchewan, Saskatoon, Saskatchewan S7N 5C6

E-mail: steelet@sask.usask.ca

Padé-improvement of four-loop β -functions in massive ϕ^4 scalar field theory is shown to predict the known five-loop contribution with astonishing (0.2%) accuracy, supporting the applicability of Padé-summations for approximating all-orders $\overline{\text{MS}}$ QCD β -functions, as suggested by Ellis, Karliner, and Samuel. Surprisingly, the most general set of [2|2] approximants consistent with known two-, three-, and four-loop contributions to the QCD β -function with up to six flavours fail to exhibit any zeros that could be interpreted as positive infrared fixed points, *regardless* of the unknown five-loop term. When they occur, positive zeros of such [2|2] approximants are preceded by singularities, leading to a double-valued β -function that is decoupled entirely from the infrared region, similar to the β -function of SUSY gluodynamics.

Higher order terms of the QCD $\overline{\text{MS}}$ β -function

$$\mu^2 \frac{dx}{d\mu^2} \equiv \beta(x), \quad (1a)$$

$$\beta(x) = - \sum_{i=0}^{\infty} \beta_i x^{i+2}, \quad (1b)$$

$x \equiv \alpha_s(\mu)/\pi$ are, upon truncation, known to permit the occurrence of fixed points other than the ultraviolet fixed point at $x = 0$; e.g. the positive infrared fixed point (IRFP) which occurs for $9 \leq n_f \leq 16$ when the series for $\beta(x)$ in (1) is truncated after two terms [$\beta_0 = (11 - 2n_f/3)/4$; $\beta_1 = (102 - 38n_f/3)/16$; $x_{IRFP} = -\beta_0/\beta_1$]. However, the fixed points arising from such truncation are likely to be spurious, as the candidate-value for x_{IRFP} is sufficiently large for the highest-order term in the series $\beta(x_{IRFP})$ to be comparable in magnitude to lower terms [e.g. $|\beta_1 x^3| = |\beta_0 x^2|$]. In a recent paper,¹ Ellis, Karliner and Samuel predicted the coefficient β_3 via Padé approximant methods, and claimed that β_{0-2} and their prediction for β_3 yield a Padé summation of the β -function with a nonzero IRFP consistent with an earlier prediction by Mattingly and Stevenson.² This Mattingly-Stevenson scenario leads to the freezing-out of the coupling to a constant value in the infrared region, as shown schematically in Fig. 1.

Padé summation of the $\overline{\text{MS}}$ β -function identifies the infinite series

$$\beta(x) = -\beta_0 x^2 (1 + R_1 x + R_2 x^2 + R_3 x^3 + \dots) \quad (2)$$

($R_i \equiv \beta_i/\beta_0$) with a Padé approximant which incorporates the known UV asymptotics of the β -function,

$$\beta(x) \rightarrow \beta_{[N|M]}(x) \equiv -\beta_0 x^2 S_{[N|M]}(x), \quad (3a)$$




Figure 1: Mattingly-Stevenson scenario

whose Maclaurin expansion reproduces the known terms in the infinite series (2):

$$\begin{aligned} S_{[N|M]}(x) &= \frac{1 + a_1 x + \dots + a_N x^N}{1 + b_1 x + \dots + b_M x^M} \\ &= 1 + R_1 x + R_2 x^2 + R_3 x^3 + \dots \quad (3b) \end{aligned}$$

An IRFP of the β -function would, in this approximation, necessarily be identified with a positive zero of $\beta_{[N|M]}$; i.e. a positive zero (x_{num}) of $1 + a_1 x + \dots + a_N x^N$, the numerator of $S_{[N|M]}$, *provided* $S_{[N|M]}$ remains positive for $0 \leq x \leq x_{num}$. This latter requirement precludes the existence of a positive zero (x_{den}) of the denominator $1 + b_1 x + \dots + b_M x^M$ that falls in the interval $0 \leq x \leq x_{num}$.

One cannot automatically dismiss the possibility of such a denominator zero occurring within the true QCD

β -function. The β -function of SU(N) SUSY gluodynamics is known *exactly* if no matter fields are present,³ and exhibits precisely such a zero:

$$\beta(x) = -\frac{3Nx^2}{4} \left[\frac{1}{1 - Nx/2} \right]. \quad (4)$$

The β -function (4) has been discussed further by Kogan and Shifman.⁴ If (4) is incorporated into (1a), the resulting Kogan-Shifman scenario (Fig. 2) for $x(\mu)$ is indicative of both a strong phase in the ultraviolet region (the upper branch of Fig. 2) as well as the existence of an infrared cut off (μ_c) on the domain of $x(\mu)$ that renders the infrared region $\mu < \mu_c$ inaccessible.^a




Figure 2: The Kogan-Shifman scenario [$x_{den} = x(\mu_c)$]

Without use of Padé summation methods, as described above, the known terms of the infinite series representation (2) for the $\overline{\text{MS}}$ QCD β -function offer little information as to whether the Mattingly-Stevenson (Fig. 1) or Kogan-Shifman (Fig. 2) scenario is appropriate for the evolution of the strong coupling. However, there is reason to believe that Padé summation representations of the β -function [eq. (3a)] may indeed be appropriate for quantum field theoretical calculations. Ellis, Gardi, Karliner, and Samuel^{1,5} have argued that Padé-summations (3) converge to their perturbative series (2) as N and M increase for any such series dominated by a finite set of renormalon poles, consistent with the following asymptotic error formula for the difference between R_{N+M+1} in (2) and the value $R_{N+M+1}^{\text{Padé}}$ predicted via use of the $[N/M]$ approximant in (3b):^{1,6}

$$\delta_{N+M+1} \equiv \frac{R_{N+M+1}^{\text{Padé}} - R_{N+M+1}}{R_{N+M+1}}$$

^aWe are assuming α_s to be real. The possibility of α_s being complex for $\mu < \mu_c$ is addressed in ref. 4.

$$= -\frac{M!A^M}{[N + M(1 + a) + b]^M}. \quad (5)$$

In (5), $\{a, b, A\}$ are constants to be determined.

To demonstrate the utility of this asymptotic error formula, consider the known β -function for massive ϕ^4 scalar-field theory:⁷

$$\mathcal{L} = \frac{1}{2}(\partial_\mu \phi)(\partial^\mu \phi) + \frac{1}{2}m^2\phi^2 + g \left(\frac{16\pi^2}{4!} \right) (\phi^2)^2, \quad (6)$$

$$\beta(g) = 1.5g^2 \left\{ 1 - \frac{17}{9}g + 10.8499g^2 - 90.5353g^3 + R_4g^4 + \dots \right\}. \quad (7)$$

Using the first two terms of (7) to generate a $[0/1]$ approximant, as in (3), one would predict $R_2^{\text{Padé}} = (-17/9)^2$, in which case we see from (5) that

$$\delta_2 = \frac{(-17/9)^2 - 10.8499}{10.8499} = \frac{-A}{1 + (a + b)}. \quad (8)$$

Using the first three terms of (7) to generate a $[1/1]$ approximant, one would predict $R_4^{\text{Padé}} = (10.8499)^2/(-17/9)$, in which case

$$\begin{aligned} \delta_3 &= \frac{[(10.8499)^2/(-17/9)] - (-90.5353)}{(-90.5353)} \\ &= \frac{-A}{2 + (a + b)}. \end{aligned} \quad (9)$$

Equations (8) and (9) are two equations for the unknown constants A and $(a + b)$, with solutions

$$A = [1/\delta_2 - 1/\delta_3]^{-1}, \quad (a + b) = (\delta_2 - 2\delta_3)/(\delta_3 - \delta_2). \quad (10)$$

We can substitute (10) into (5) to determine R_4 . The first three terms in the series (7) generate a $[2/1]$ approximant whose Maclaurin expansion (3b) predicts $R_4^{\text{Padé}} = (-90.5353)^2/(10.8499)$. Upon substitution of A , $(a + b)$, and $R_4^{\text{Padé}}$ into (5) we find that this asymptotic error formula *predicts* that

$$\begin{aligned} R_4 &= R_4^{\text{Padé}}/(1 + \delta_4) \\ &= R_4^{\text{Padé}}[3 + (a + b)]/[3 + (a + b) - A] = 947.8 \end{aligned} \quad (11)$$

The value of R_4 in (7) has been explicitly calculated⁷ to be 949.5, in very close agreement with (11). The series of steps leading from (5) to (11), a methodological recipe first presented in ref. 6, has also been applied^{1,8} to N -component scalar field theory, for which the Lagrangian (6) is modified such that $\phi \rightarrow \phi_a$, $\phi^2 \rightarrow \sum_{a=1}^N \phi_a^2$, $N = \{2, 3, 4\}$. Agreement with calculated values of R_4 ($R_4 \equiv \beta_4/\beta_0$) remains within 3.5% for $N \leq 4$.⁸

This startling agreement suggests that Padé methodology may also be applicable to the QCD β -function, particularly in the $n_f = 0$ gluodynamic limit where

1. such methods are expected to be most accurate,⁶
2. comparison with the Kogan-Shifman scenario for SUSY gluodynamics is most relevant.

For $n_f = 0$, the 4-loop $\overline{\text{MS}}$ QCD β -function, as defined by (1a), is given by⁹

$$\beta(x) = -\frac{11}{4}x^2[1 + 2.31818 + 8.11648x^2 + 41.5383x^3 + \sum_{k=4}^{\infty} R_k x^k]. \quad (12)$$

The coefficients R_k are presently not known for $k \geq 4$. The first three terms in the series (12) are sufficient in themselves to determine the Padé approximants $S_{[1|2]}$ and $S_{[2|1]}$, as defined in (3). These approximants are

$$\beta_{[2|1]}(x) = -\frac{11}{4}x^2 \left[\frac{1 - 2.7996x - 3.7475x^2}{1 - 5.1178x} \right], \quad (13)$$

$$\beta_{[1|2]}(x) = -\frac{11}{4}x^2 \left[\frac{1 - 5.9672x}{1 - 8.2854x + 11.091x^2} \right]. \quad (14)$$

In both (13) and (14), the (first) positive denominator zero *precedes* the positive numerator zero: for (13), $x_{num} = 0.264 > x_{den} = 0.195$; for (14), $x_{num} = 0.168 > x_{den} = 0.151$. Consequently, x_{num} cannot be identified with the Mattingly-Stevenson IRFP in either case, as this zero is separated from the small x -region by a singularity past which the β -function switches sign. Indeed the ordering $0 < x_{den} < x_{num}$ is suggestive of a Kogan-Shifman scenario in which x_{num} , if taken seriously, is an *ultraviolet* fixed point (UVFP) characterizing the strong phase [*i.e.*, the upper branch of Fig 2].

We can apply the asymptotic error formula (5) to the series (12) in precisely the same way we applied it to (7). We then obtain an estimate $R_4 = 302.2$, analogous to (11). Using this value of R_4 in conjunction with the known terms of (12), it is possible to obtain a $[2|2]$ -approximant β -function

$$\beta_{[2|2]}(x) = -\frac{11}{4}x^2 \left[\frac{1 - 9.6296x + 4.3327x^2}{1 - 11.9477x + 23.913x^2} \right]. \quad (15)$$

The first positive numerator zero $x_{num} = 0.1092$ is again larger than the first positive denominator zero $x_{den} = 0.1063$, precluding the identification of x_{num} as the IRFP of the Mattingly-Stevenson scenario (Fig. 1). Instead, the β -function (15) is consistent with the Kogan-Shifman scenario of Fig. 2, with x_{num} again identified as a nonzero UVFP for the strong phase.

Curiously, the ordering $0 < x_{den} < x_{num}$ characterizes $[2|2]$ -approximant β -functions *even if* R_4 is allowed to be arbitrary. The most general such β -function that

reproduces the first four terms of (12) [the first three being known] is

$$\beta_{[2|2]}(x) = -\frac{11}{4}x^2 \times \left[\frac{1 + (13.403 - 0.076215R_4)x - (22.915 - 0.090166R_4)x^2}{1 + (11.084 - 0.076215R_4)x - (56.727 - 0.26685R_4)x^2} \right]. \quad (16)$$

It is easy to verify the first positive numerator zero of (16) is always larger than the first positive denominator zero [Fig. 3], although these zeros become asymptotically close as $R_4 \rightarrow +\infty$. Thus, we see that the first positive zero of *any* $[2|2]$ Padé approximant whose Maclaurin expansion reproduces the known terms of eq. (12) *cannot* be identified as an IRFP, nor is such Padé-summation indicative of a Fig. 1 scenario for the $\overline{\text{MS}}$ $n_f = 0$ β -function.

Remarkably, the same set of conclusions can be drawn for the physically interesting case of three light flavours. When $n_f = 3$, the 4-loop $\overline{\text{MS}}$ QCD β -function is given by⁹

$$\beta(x) = -\frac{9x^2}{4} \left[1 + (16/9)x + 4.471065x^2 + 20.99027x^3 + \sum_{k=4}^{\infty} R_k x^k \right], \quad (17)$$

with R_k not presently known for $k \geq 4$. The known terms in (17) determine $[2|1]$ and $[1|2]$ Padé-summation representations of the $n_f = 3$ β -function,

$$\beta_{[2|1]}(x) = -\frac{9x^2}{4} \left[\frac{1 - 2.91691x - 3.87504x^2}{1 - 4.69468x} \right], \quad (18)$$

$$\beta_{[1|2]}(x) = -\frac{9x^2}{4} \left[\frac{1 - 8.17337x}{1 - 9.95115x + 13.2199x^2} \right]. \quad (19)$$

The positive zero of $\beta_{[2|1]}(x)$ ($x = 0.2559$) occurs after the pole at $x = 0.2130$; the positive zero of $\beta_{[2|1]}(x)$ at $x = 0.1223$ similarly occurs after a pole at $x = 0.1194$. The most general $[2|2]$ approximant consistent with (17) is

$$\beta_{[2|2]}(x) = -\frac{9x^2}{4} \times \left[\frac{1 + (7.1945 - 0.10261R_4)x - (11.329 - 0.075643R_4)x^2}{1 + (5.4168 - 0.10261R_4)x - (25.430 - 0.25806R_4)x^2} \right]. \quad (20)$$

The Maclaurin expansion of (20) reproduces the series in (17), including its (unknown) $R_4 x^4$ term. As was the case in (16), the first positive zero of the denominator of (20) is always seen to precede the first positive zero of the numerator, regardless of R_4 . Thus the $[1|2]$, $[2|1]$ and most general possible $[2|2]$ -approximant representations of the $n_f = 3$ $\overline{\text{MS}}$ β -function uphold the ordering

$0 < x_{den} < x_{num}$, an ordering that precludes the identification of x_{num} with the IRFP of the Mattingly-Stevenson scenario. Moreover, [2|2]-approximant β -functions for arbitrary R_4 have been constructed⁸ analogous to (16) and (20) for $n_f = \{4, 5, 6\}$, and for each of these, the $0 < x_{den} < x_{num}$ ordering persists regardless of R_4 . A range for R_4 for which an ordering compatible with Fig. 1 ($0 < x_{num} < x_{den}$) is possible does not occur until $n_f = 7$.

As noted above, the ordering $0 < x_{den} < x_{num}$ suggests the occurrence of a double-valued QCD coupling constant, as is the case in SUSY gluodynamics (Fig. 2). Such a scenario is seen to decouple the infrared region $\mu < \mu_c$ from the domain of α_s , provided α_s is understood to be real. Such a scenario is also indicative of a strong phase at short distances⁴ with possible implications for dynamical electroweak symmetry breaking, suggesting that QCD may even furnish its own “technicolour.”




Figure 3: Dependence of the first positive numerator zero (x_{num}) and denominator zero (x_{den}) of (16) on R_4 , the (presently unknown) five-loop coefficient of the $n_f = 0$ β -function.

Acknowledgements

VE and TGS are grateful for research support from NSERC, the Natural Sciences and Engineering Research Council of Canada.

References

1. J. Ellis, M. Karliner and M.A. Samuel, *Phys. Rev. D* **400**, 176 (1997).
2. A.C. Mattingly and P. M. Stevenson, *Phys. Rev. D* **49**, 437 (1994), and *Phys. Rev. Lett.* **69**, 1320 (1992).
3. V. Novikov, M. Shifman, A. Vainshtein, and V. Zakharov, *Nucl. Phys. B* **229**, 381 (1983), and *Phys. Lett. B* **166**, 329 (1986); D.R.T. Jones, *Phys. Lett. B* **123**, 45 (1983).
4. I. I. Kogan and M. Shifman, *Phys. Rev. Lett.* **75**, 2085 (1995).
5. M. Samuel, J. Ellis, and M. Karliner, *Phys. Rev. Lett.* **74**, 4380 (1995); J. Ellis, E. Gardi, M. Karliner, and M. A. Samuel, *Phys. Lett. B* **366**, 268 (1996) and *Phys. Rev. D* **54**, 6986 (1996).
6. J. Ellis, I. Jack, D.R.T. Jones, M. Karliner, and M. A. Samuel, *Phys. Rev. D* **57**, 2665 (1998).
7. H. Kleinert, J. Neu, V. Schulte-Frohlinde, K.G. Chetyrkin, and S. A. Larin, *Phys. Lett. B* **272**, 39 (1991).
8. V. Elias, T. G. Steele, F. Chishtie, R. Migneron, and K. Sprague, *Phys. Rev. D*, (to appear) hep-ph/9806324.
9. T. van Ritbergen, J.A.M. Vermaseren, and S.A. Larin, *Phys. Lett. B* **400**, 379 (1997).